



# Arkansas Audubon Society

## 50th Anniversary Field List

Date \_\_\_\_\_ Time \_\_\_\_\_ Total Species Observed \_\_\_\_\_

Location \_\_\_\_\_ County \_\_\_\_\_

Observers \_\_\_\_\_

This field list provides generalized information on 396 species of birds that have occurred in Arkansas. It is based primarily on the card file maintained by the Arkansas Audubon Society, and secondarily on other sources, including *Arkansas Birds-Their Distribution and Abundance* (James and Neal, UA Press, 1986), *Breeding Bird Surveys*, Christmas Bird Counts, and published literature on Arkansas birds. Since the information that follows is generalized, it may not seem appropriate for a given species in some locales. Taxonomy and nomenclature follow the *Checklist of North American Birds*, Seventh Edition, AOU, 1998, as amended through the Forty-fifth supplement, July 2004.

Approximate dates of arrival and departure are indicated by numbers 1-12 for months of the year and letters A, B, C, D for the weeks of the month. A plus (+) sign immediately following indicates that discontinuous records outside this period are on file. For species that have nested in Arkansas, one of two symbols follows the name: an asterisk (\*) means that the species currently nests with regularity, either yearly, or at least, with some consistency; an open circle (°) means that the species does not nest regularly and that there are few or no such reports since 1950. Beneath the dates of arrival and departure are codes which categorize seasonal status, relative abundance, and, in some cases, the sections of the state of primary occurrence.

The months of the year are grouped into the following seasons:

Sp-spring (March-May) S-summer (June-July)  
F-fall (August-November) W-winter (December-February)\*

In combination with these designations, the following abbreviations are used to denote seasonal status:

P-Present, species seen in the state each month but not necessarily in all sections (Great Egret)  
M-Migrant, species passes through the state in the spring and/or fall (Cape May Warbler)  
R-Resident, species remains for an extended period of time (Painted Bunting)  
L-Local, species occurs in only a few places (Great-tailed Grackle)  
V-Visitor, species usually not present for an extended time (Black-legged Kittiwake)

The frequencies of occurrence assume reasonable skill spent in suitable habitat during appropriate seasons and are defined as follows:

c-common, always or almost always observed (Killdeer)  
fc-fairly common, seen on a majority of trips but usually in smaller numbers (Red-shouldered Hawk)  
u-uncommon, may be difficult to find; seen only on a minority of trips (Golden-winged Warbler)  
r-rare, may be seen annually or nearly so but not in all locales (Golden Eagle)  
vr-very rare, encountered at irregular or infrequent intervals (Ruff)  
occ-occasional, 3-4 state records overall with at least 2 records since 1950 (Long-tailed Jaeger)  
acc-accidental, 1-2 state records (Fork-tailed Flycatcher)  
irr-irruptive, variable in occurrence, number, and distribution (Red-breasted Nuthatch)

The abbreviations in parentheses following the statements of relative abundance indicate the primary but not necessarily the exclusive section of the state where the species occurs: (n) north, (s) south, (e) east, (w) west, (c) central, (nc) north-central, (sc) south-central, (ec) east-central, (wc) west-central, (ne) northeast, (se) southeast, (nw) northwest, and (sw) southwest.

### Reporting Bird Sightings

Ornithology has benefited a great deal from the contributions of amateurs over the years. Birders who faithfully report significant observations and make them a part of the permanent record help ornithologists better understand the intricacies of bird distribution and abundance. Although it isn't always obvious whether a sighting should be reported, a good rule to follow is "If in doubt, make the report." More specific instances in which a report should be made include:

- If a species appears outside of its normal season or geographical region, it should be reported.
- If a large number of a certain species is encountered, that count of individuals should be reported.
- If an unusual or unlisted species is seen, report it by all means but keep in mind that the greater the rarity the more thorough supporting documentation should be.

The basic information needed for all reports includes: common name, age (adult, immature, etc.), plumage (breeding, non-breeding, etc.), number seen, location (county, nearest town, etc.), and date. The more unusual reports require additional information such as time seen, light conditions, optical equipment, distance to bird and how measured, as well as a description of the bird's size, shape, color pattern, vocalizations if any were heard, and behavior. It is also important to eliminate any similarly appearing species and discuss prior experience with the species being reported and any with it might be confused with. The names and addresses of additional observers should be included as well as any reference material consulted and how it influenced the description. If it is possible to obtain photographs or audio recordings, be sure to include copies with the report. **Above all else, remember that the documentation process is in place to gather additional, substantiating information and is not meant to intimidate potential contributors or question the authenticity of their reports.**

Reports of birds should be sent to Max Parker, 14300 Chenal Parkway #7473, Little Rock AR, 72211 or submitted online via the society's web-site at [www.arbirds.org](http://www.arbirds.org). **Written and/or photographic documentation is required when reporting rarities or species observed out of season.** The Arkansas Bird Records Committee reviews those reports.

The Arkansas Audubon Society was formed in May 1955 to promote the protection and wise use of our natural resources. Weekend meetings are held twice each year at locations throughout the state and are generally scheduled to coincide with the peaks of spring and fall migration. Guest speakers, workshops, and guided field trips are featured at all meetings. A quarterly newsletter is published to keep the membership informed regarding current conservation issues, noteworthy bird records, and news of members and upcoming events. Membership is open to all interested persons. For further information, please visit the society's website at [www.arbirds.org](http://www.arbirds.org)



The 50th anniversary checklist was compiled by Max Parker. Document design and typesetting by Charles Mills and Dolores Harrington. Editorial assistance provided by Leif Anderson, Douglas A. James, Norman Lavers, Cheryl Lavers, Bill Lisowsky, Mike Mlodnow, Joe Neal, Kenny Nichols, LaDonna Nichols and Helen Parker. Graphics provided by Sharon Carter and Pat Moore.

Price: 50¢ each 10 copies: \$4.50 25 copies: \$10 50 copies: \$15

Prices quoted for quantity purchases do not include postage.

For one copy of this field list, send 50¢ and a long self-addressed, self-stamped envelope to Max Parker, 14300 Chenal Parkway #7473, Little Rock AR, 72211.

MC389EX7 (2/28/05)

<b>Order Anseriformes</b> <b>Whistling-Ducks, Swans, Geese, and Ducks</b>	Scoter, Surf	10B-1D FMr SpMvr	
Whistling-Duck, Black-bellied*	3A-11D+ SRLu WVvr	White-winged	10D-5D+ WVvr
Fulvous°	4C-5C 7C-9D SpVlr SVLr	Black	10D-3C WVr
Goose, Greater White-fronted	10A-3C+ Mfc WRLfc	Duck, Long-tailed	11A-4A+ WVu
Snow	10A-4C+ Mc WRc (c, ce)	Bufflehead	11B-4C+ WRLu
Ross's	10D-4C Mfc WRLfc	Goldeneye, Common	11B-4C WRLu
Brant	acc (1, 1990)	Barrow's	acc (2, 2003)
Goose, Cackling	Status Under Review Fully document all sightings	Merganser, Hooded*	P SRLu WRLfc
Canada	9A-4D+ Mc WRc (c, ce)	Common	10D-3C+ WVr
Swan, Mute	11D-2D+ WVvr	Red-breasted	10D-5B+ WRu
Trumpeter	11A-3A WRLu	Duck, Ruddy°	10A-4C+ WRc SRr
Tundra	11C-3D Mr WVr (c, ce)	<b>Order Galliformes</b> <b>Grouse, Turkey, and Quail</b>	
Duck, Wood*	P SRc WRLc	Grouse, Ruffed*	P Lvr (nw, nc)
Gadwall	10B-4C+ WRc	Turkey, Wild*	P fc
Wigeon, American	9C-4C+ Mc WRc	Bobwhite, Northern*	P Lfc
Duck, American Black	10A-3B+ WRLu (c)	<b>Order Gaviiformes</b> <b>Loons</b>	
Mallard*	P SRu	Loon, Red-throated	10D-4A WVvr
Duck, Mottled	4B-8C SRr (se, ce, sw)	Pacific	11B-5D WVvr
Teal, Blue-winged*	P Mc SRr WRr	Common	10A-5D+ Mfc WRLfc (ce, s)
Cinnamon	2C-5B 9C-11D+ Mvr (w)	Yellow-billed	acc (1, 1991)
Shoveler, Northern	9A-4C+ Mc WRc (c, sw)	<b>Order Podicipediformes</b> <b>Grebes</b>	
Pintail, Northern	10A-4B+ Mc WRc (c, sw)	Grebe, Pied-billed*	P WRc SRLu
Garganey	acc (1, 1984)	Horned	10A-3D+ WRLfc
Teal, Green-winged	10A-4C+ WRc	Red-necked	occ (4, 2004)
Canvasback	10D-4D+ WRLfc	Eared	9A-5A+ Mfc WRu
Redhead	10D-4A+ WRu	Western	10B-3D+ WVr
Duck, Ring-necked	10C-5B+ WRc	<b>Order Pelecaniformes</b> <b>Pelicans, Cormorants, and Darters</b>	
Tufted	acc (1, 2001)	Pelican, American White	P Mfc SVu WRLfc
Scaup, Greater	10A-4B Mu WRu	Brown	3C-11B+ Vr
Lesser	10D-4D+ Mc WRc	Cormorant, Neotropic°	3D-12C SRvr

Cormorant, Double-crested*	P Mc SRLu WRLc	Hawk, Cooper's*	P WRu SRr
Anhinga*	3C-12C+ SRLu WVvr (s)	Goshawk, Northern	11A-1B WVvr
Frigatebird, Magnificent	9B-10D+ FVvr	Hawk, Red-shouldered*	P fc
<b>Order Ciconiiformes</b> <b>Herons, Egrets, Ibises, and Vultures</b>		Broad-winged*	3D-10A+ Mfc SRu
Bittern, American°	3A-5D 9B-12D+ Mu WVvr	Swainson's*	3D-5D 9C-11C+ SRu (nw)
Least*	4A-10A+ SRLr	Red-tailed*	P c
Heron, Great Blue*	P c	Ferruginous	10D-4B WVvr
Egret, Great*	P SRLc WRLfc (ce, s)	Rough-legged	10D-3C WVr
Snowy*	3C-11C+ Mfc SRLfc	Eagle, Golden	9B-3B+ WRLr
Heron, Little Blue*	3C-10D+ Mc SRLc	Kestrel, American*	P Mc SRu WRc
Tricolored°	4A-11A+ SRr (s) FVu (s)	Merlin	9B-4C+ Mu WVu
Egret, Cattle*	3B-11B+ Mc SRc WVr (s)	Falcon, Peregrine°	9B-5B+ Mu WVr
Heron, Green*	3B-10D+ SRc	Prairie	11B-2B WVvr
Night-Heron, Black-crowned*	3B-11A+ Mu SRLu	<b>Order Gruiformes</b> <b>Rails, Gallinules, and Cranes</b>	
Yellow-crowned*	4B-10C+ SRLu	Rail, Yellow	9A-5B+ Mr WVvr
Ibis, White*	3D-11A+ SRLc (s, sw)	Black	9B-10B FMvr
Glossy°	4C-5D 8C-10D+ Mr	King*	P SRLr WRvr
White-faced	4B-5D 8D-9D+ Mr	Virginia	4B-5D 8B-12D+ MLu WRLvr
Spoonbill, Roseate	7B-10B FVr (s)	Sora	4A-5B 9A-12D+ Mu WRr (ce, s)
Stork, Wood	6B-11A FVu (s, e)	Gallinule, Purple*	4C-10C SRLr (ce, s)
Vulture, Black*	P u (n) fc (ce, s)	Moorhen, Common*	4C-12D+ SRLu (ce, s) WRLr (sw)
Turkey*	P c	Coot, American*	P WRc SRLu
<b>Order Falconiformes</b> <b>Hawks and Falcons</b>		Crane, Sandhill	9D-5C WMr
Osprey*	4A-5D 8C-11D+ Mc SRLr WRr	<b>Order Charadriiformes</b> <b>Plovers, Sandpipers, Gulls, and Terns</b>	
Kite, Swallow-tailed°	4A-9D SRLvr	Plover, Black-bellied	3C-5D 8D-12D+ SpMu FMu
White-tailed	3A-4D+ SpVvr	Golden-Plover, Am.	3A-5D 9C-12A+ SpMc FMu
Mississippi*	4C-9C+ SRLfc	Plover, Snowy	occ (3, 2001)
Eagle, Bald*	WRfc SRLu	Wilson's	occ (3, 1998)
Harrier, Northern°	8C-4B+ WRfc (u, nw)	Semipalmated	4C-5C 7B-9D+ Mc
Hawk, Sharp-shinned*	P Mfc WRu SRr (n)	Piping	4B-5A 7B-10B SpMr FMr

Killdeer*	P c	Dowitcher, Long-billed	3D-5B 7A-12D+ SpMfc FMfc	
Plover, Mountain	acc (1, 1951)	Snipe, Wilson's	8D-5B+ Mc WRc (s)	
Stilt, Black-necked*	4A-10D SRfc (ec, se)	Woodcock, American*	P u	
Avocet, American	4B-5C 7A-11C+ SpMr FMu	Phalarope, Wilson's	3C-5D 7A-10D Mfc	
Yellowlegs, Greater	3B-5B 7A-12D+ Mfc WRu	Red-necked	5C-6A 7B-11B Mr	
Lesser	2B-5D 6D-12D+ Mc WRu	Red	8D-11D+ FMvr	
Sandpiper, Solitary	3D-5B 6B-12D+ Mfc	Jaeger, Pomarine	acc (1, 1983)	
Willet	3D-5D 6C-9D+ SpMu FMu	Parasitic	9A-12B FVvr	
Sandpiper, Spotted°	3D-6B 7A-12D+ Mc WVr	Long-tailed	occ (4, 2000)	
Upland	3D-5C 6D-10D+ Mu	Gull, Laughing	8A-7B Vu	
Whimbrel	5B-5D 9A-9B Mr	Franklin's	3C-6A 9D-12C+ SpMr FMfc	
Curlew, Long-billed	10B-11C+ FMvr	Little	11D-3C+ WVvr	
Godwit, Hudsonian	4B-5D 8C-10B SpMu FMr	Black-headed	11C-1B WVvr	
Marbled	4B-5D 8B-10B SpMvr FMr	Bonaparte's	9D-4D+ Mfc WRfc	
Turnstone, Ruddy	5B-6B 7D-10B Mr	Ring-billed	8C-5A+ WRc	
Knot, Red	8C-10A FMvr	California	acc (1, 2004)	
Sanderling	4C-5D 7C-10D+ SpMu FMfc	Herring	8B-5C+ WRLu	
Sandpiper, Semipalmated	4A-6B 7B-11A+ Mc	Thayer's	acc (2, 2005)	
Western	4B-6A 7A-12D+ SpMu FMfc	Lesser Black-backed	11C-2C WVvr	
Least	7B-6B+ Mc WRu	Glaucous	11D-2A WVvr	
White-rumped	4B-6C 8B-10D SpMLfc FMvr	Sabine's	9C-10B+ FVLr	
Baird's	3B-6A 7D-11C SpMu FMfc	Kittiwake, Black-legged	11B-3A WVvr	
Pectoral	3A-6B 6D-11D+ Mc	Tern, Caspian	4A-6C 7A-10C+ SpMu FMfc	
Purple	acc (1, 1976)	Royal	occ (4, 2002)	
Dunlin	4B-5D 10A-12D+ Mfc WRu	Common	4D-5D 7A-10D SpMr FMu	
Sandpiper, Curlew	acc (2, 2004)	Forster's	4B-2D+ Mfc WRLu (s)	
Stilt	3D-6A 7A-11D+ SpMu FMfc	Least*	5A-9D+ Mu SRLu	
Buff-breasted	4A-5C 7D-10A+ SpMr FMfc	Bridled	acc (1, 1985)	
Ruff	3D-5B+ SpMvr	Sooty	occ (3, 2002)	
Dowitcher, Short-billed	4D-6A 7D-10A SpMu FMu	Black	4B-6B 6D-10C+ Mc	

Skimmer, Black	occ (3, 1969)	Swift, White-throated	acc (3, 1999)	
Murrelet, Long-billed	acc (1, 1980)	Violet-ear, Green	6A-10A FVvr	
<b>Order Columbiformes Pigeons and Doves</b>		Hummingbird, Buff-bellied	acc (1, 1993)	
Pigeon, Rock*	P c	Magnificent	acc (1, 1993)	
Collared-Dove, Eurasian*	P Lc	Ruby-throated*	3B-11B+ SRc	
Dove, White-winged	11C-6D Vu	Black-chinned	10D-3D WVvr	
Mourning*	P c	Anna's	10C-3B WVvr	
Inca*	P Lfc	Calliope	occ (4, 2004)	
Ground-Dove, Common	8C-5A Vr	Broad-tailed	occ (3, 2003)	
<b>Order Cuculiformes Cuckoos, Roadrunners, and Anis</b>		Rufous	8B-4A FMr WVr	
Cuckoo, Black-billed°	4D-5D 8A-10C+ Mu SRvr (n)	<b>Order Coraciiformes Kingfishers</b>		
Yellow-billed*	4A-10C+ SRc	Kingfisher, Belted*	P c	
Roadrunner, Greater*	P Lu (w, ce)	<b>Order Piciformes Woodpeckers</b>		
Ani, Groove-billed	9C-11D+ SpVvr FVr	Woodpecker, Lewis's	acc (1, 1967)	
<b>Order Strigiformes Owls</b>		Red-headed*	P fc	
Owl, Barn*	P Lu	Red-bellied*	P c	
Screech-Owl, Eastern*	P fc	Sapsucker, Yellow-bellied	9A-5A+ WRc	
Owl, Great Horned*	P fc	Woodpecker, Downy*	P c	
Snowy	acc (4, 1955)	Hairy*	P fc	
Burrowing	9D-3D WVLvr (e, ce)	Red-cockaded*	P Lr (ec, sc, wc)	
Barred*	P fc	Flicker, Northern*	P SRu WRc	
Long-eared	11A-5D+ WRr	Woodpecker, Pileated*	P fc	
Short-eared	10D-4D WRLu	<b>Order Passeriformes Family Tyrannidae Flycatchers</b>		
Northern Saw-whet	11A-2D WVvr	Flycatcher, Olive-sided	4D-6D 8A-10B Mfc	
<b>Order Caprimulgiformes Goatsuckers</b>		Wood-Pewee, Eastern*	4B-10C SRc	
Nighthawk, Common*	4B-11A+ SRc	Flycatcher, Yellow-bellied	4D-5D 8B-10C Mr	
Chuck-wills-widow*	3D-10B SRc	Acadian*	4B-9C SRfc	
Whip-poor-will*	3D-10B SRfc (n, ce, wc)	Alder	5A-6A 8A-9B Mu	
<b>Order Apodiformes Swifts and Hummingbirds</b>		Willow*	5A-9C SpMu SRLr (nw)	
Swift, Chimney*	3B-11A SRc	Least	4D-5C 8D-9D+ SpMfc FMu	

Phoebe, Eastern*	P SRc (n, wc) WRfc (s)	Swallow, N. Rough-winged*	3C-10B+ SRc
Say's	4A-4D 9C-12D SpVvr FVvr	Bank*	4A-5D 7A-9D+ Mfc SRvr
Flycatcher, Vermilion	10A-4B+ WVvr	Cliff*	3D-10C+ Mc SRc
Great Crested*	4B-10A+ SRc	Barn*	3A-11D+ SRc
Kingbird, Couch's	acc (1, 1995)	<b>Family Paridae</b> <b>Chickadees and Titmice</b>	
Cassin's	acc (1, 1967)	Chickadee, Carolina*	P c
Western*	3D-10C SRLu (w)	Titmouse, Tufted*	P c
Eastern*	3D-10D SRc	<b>Family Sittidae</b> <b>Nuthatches</b>	
Flycatcher, Scissor-tailed*	3D-11D+ SRc	Nuthatch, Red-breasted	9B-5B+ WRrr
Fork-tailed	acc (1, 1995)	White-breasted*	P c (n, wc)
<b>Family Laniidae</b> <b>Shrikes</b>		Brown-headed*	P c (s, wc)
Shrike, Loggerhead*	P c (c) u (w)	<b>Family Certhiidae</b> <b>Creepers</b>	
<b>Family Vireonidae</b> <b>Vireos</b>		Creeper, Brown	10A-4C+ WRfc
Vireo, White-eyed*	3C-10B+ SRc	<b>Family Troglodytidae</b> <b>Wrens</b>	
Bell's*	4C-9B+ Mu SRu	Wren, Rock	10B-3C+ WRr (w)
Yellow-throated*	3D-10A+ SRfc	Carolina*	P c
Blue-headed	9A-5D+ Mfc WRu (s)	Bewick's*	P r
Warbling*	4C-9D+ SRLu	House*	P Mfc WRu SRu (nw)
Philadelphia	4D-5D 9B-9D+ Mfc	Winter	10A-4D+ WRfc
Red-eyed*	4B-10B+ SRc	Sedge°	P SRvr (c, nw) Mfc WRu (s)
<b>Family Corvidae</b> <b>Jays and Crows</b>		Marsh	9C-5C Mfc WRu
Jay, Blue*	P c	<b>Family Regulidae</b> <b>Kinglets</b>	
Nutcracker, Clark's	acc (1, 1891)	Kinglet, Golden-crowned	9D-4D WRfc
Crow, American*	P c	Ruby-crowned	9D-4D+ WRc (s, ce)
Fish*	P SRc WRLfc (cc) WRc (s)	<b>Family Sylviidae</b> <b>Gnatcatchers</b>	
<b>Family Alaudidae</b> <b>Larks</b>		Gnatcatcher, Blue-gray*	3C-9C+ SRc WVr
Lark, Horned*	P WRc SRLu	<b>Family Turdidae</b> <b>Wheatears, Bluebirds, and Thrushes</b>	
<b>Family Hirundinidae</b> <b>Swallows</b>		Wheatear, Northern	acc (1, 1990)
Martin, Purple*	2A-10B+ SRc	Bluebird, Eastern*	P c
Swallow, Tree*	2C-11C+ Mc SRLfc	Mountain	acc (1, 1985)

Solitaire, Townsend's	10C-2C WVvr	Warbler, Cape May	4D-5D 8D-11D+ Mr
Veery	4C-5C+ SpMu	Black-throated Blue	5C-5D 9C-10C+ Mvr
Thrush, Gray-cheeked	4B-5D+ SpMfc	Yellow-rumped	10A-4D+ WRc
Swainson's	4A-5D 9B-10C+ SpMc FMu	Black-throated Green*	3D-10C+ Mc SRLu (nw, w)
Hermit	10B-4C+ WRc (cc, s)	Blackburnian	4D-6A 9A-10B SpMu FMu
Wood*	4B-9C+ SRfc	Yellow-throated*	3D-10C+ SRfc
Robin, American*	P c	Pine*	P c (wc, s, nw)
<b>Family Mimidae</b> <b>Mockingbirds, Thrashers, and Allies</b>		Prairie*	4B-9C+ SRfc
Catbird, Gray*	4C-10C+ SRc WRr (s)	Palm	4C-5A 9C-11C+ SpMu FMu WVr
Mockingbird, Northern*	P c	Bay-breasted	4B-6B 9D-10B+ Mu
Thrasher, Sage	acc (2, 2004)	Blackpoll	4B-5D SpMfc
Brown*	P c	Cerulean*	4C-8C+ SRu
<b>Family Sturnidae</b> <b>Starlings</b>		Black-and-white*	3C-10A+ SRc
Starling, European*	P c	Redstart, American*	4C-10C Mfc SRu
<b>Family Motacillidae</b> <b>Pipits</b>		Warbler, Prothonotary*	4A-9C+ SRfc
Pipit, American	9D-4B+ Mfc WRfc	Worm-eating*	4A-9D+ SRu (sc, nw)
Sprague's	10A-4C WRvr	Swainson's*	4A-9C+ SRLfc
<b>Family Bombycillidae</b> <b>Waxwings</b>		Ovenbird*	4C-10B+ SRfc (n, ce, wc)
Waxwing, Bohemian	1D-3C+ WVvr	Waterthrush, Northern	4C-5C 7B-9D+ Mfc
Cedar*	P SRu (nw) WRc	Louisiana*	3C-10B SRfc (w)
<b>Family Parulidae</b> <b>Wood Warblers</b>		Warbler, Kentucky*	4B-9D+ SRfc
Warbler, Blue-winged*	4B-10B Mfc SRLu (n)	Connecticut	4C-5D SpMvr
Golden-winged	4D-5B 9A-10B Mu	Mourning	5A-5D 9A-9D+ SpMu FMu
Tennessee	4C-5B 9B-10B+ Mc	Yellowthroat, Common*	4B-10C+ SRc WRr
Orange-crowned	9D-5C+ Mfc WRu (s)	Warbler, Hooded*	4A-10A+ SRfc
Nashville	4C-5B 9D-10D+ Mfc	Wilson's	4D-6A 8D-10A+ Mu
Parula, Northern*	3C-10C+ SRc	Canada	4D-6A 8D-10A Mu
Warbler, Yellow*	4C-9B+ Mfc SRLu (n)	Chat, Yellow-breasted*	4C-9D+ SRc
Chestnut-sided°	4D-10B+ SpMfc FMu SRLt (nw)	<b>Family Thrupidae</b> <b>Tanagers</b>	
Magnolia	4D-6A 9A-10B+ SpMfc FMu	Tanager, Summer*	4C-10B+ SRc WVvr (cc, s)

